

(CACFP) Infant Meal Patterns

Infants	Birth through 5 months	6 through 11 months
Breakfast, Lunch, or Supper	4-6 fluid ounces breastmilk ¹ or formula ²	6-8 fluid ounces breastmilk ¹ or formula; ² and 0-½ ounce equivalent infant cereal; ^{2,3} or 0-4 tablespoons meat, fish, poultry, whole egg, cooked dry beans, or cooked dry peas; or 0-2 ounces of cheese; or 0-4 ounces (volume) of cottage cheese; or 0-4 ounces or ½ cup of yogurt; ⁴ or a combination of the above; ⁵ and 0-2 tablespoons vegetable or fruit, or a combination of both. ^{5,6}
Snack	4-6 fluid ounces breastmilk ¹ or formula ²	2-4 fluid ounces breastmilk ¹ or formula; ² and 0-½ ounce equivalent bread; ^{3,7} or 0-¼ ounce equivalent crackers; ^{3,7} or 0-½ ounce equivalent infant cereal; ^{2,3} or 0-¼ ounce equivalent ready-to-eat breakfast cereal; ^{3,5,7,8} and 0-2 tablespoons vegetable or fruit, or a combination of both. ^{5,6}

¹ Breastmilk or formula, or portions of both, must be served; however, it is recommended that breastmilk be served in place of formula from birth through 11 months. For some breastfed infants who regularly consume less than the minimum amount of breastmilk per feeding, a serving of less than the minimum amount of breastmilk may be offered, with additional breastmilk offered at a later time if the infant will consume more.

² Infant formula and dry infant cereal must be iron-fortified.

³ Refer to FNS guidance for additional information on crediting different types of grains.

⁴ Yogurt must contain no more than 23 grams of total sugars per 6 ounces.

⁵ A serving of this component is required when the infant is developmentally ready to accept it.

⁶ Fruit and vegetable juices must not be served.

⁷ A serving of grains must be whole grain-rich, enriched meal, or enriched flour.

⁸ Breakfast cereals must contain no more than 6 grams of sugar per dry ounce (no more than 21.2 grams sucrose and other sugars per 100 grams of dry cereal).

